



## The rotating water glass

a. Consider a cube of water of volume  $\ell \times \ell \times \ell$  where  $\ell$  is “quite small”.

Top view



Side view



The forces due to pressure are denoted  $F_{t,1}$  (force tangential, 1) and so forth.

Use  $\sum \vec{F} = m\vec{a}$ :

Tangential component:

$$\begin{aligned} F_{t,1} - F_{t,2} &= ma_t = 0 \\ \implies p_{t,1}\ell^2 &= p_{t,2}\ell^2 \implies p_{t,1} = p_{t,2} \end{aligned}$$

Thus the pressure is independent of angle (which is also clear from symmetry).

Vertical component:

$$\begin{aligned} F_{v,1} - \rho V g - F_{v,2} &= ma_v = 0 \\ \implies p_{v,1}\ell^2 - \rho\ell^3 g - p_{v,2}\ell^2 &= 0 \implies p_{v,2} - p_{v,1} = -\rho\ell g \end{aligned}$$

Thus  $\frac{\partial p}{\partial h} = -\rho g$ .

Radial component:


$$\begin{aligned} F_{r,1} - F_{r,2} &= ma_r = -\rho V \omega^2 r \quad (\text{centripetal acceleration}) \\ \implies p(r)\ell^2 - p(r+\ell)\ell^2 &= -\rho\ell^3\omega^2 r \end{aligned}$$

Thus  $\frac{\partial p}{\partial r} = \rho\omega^2 r$ .

b. When moving along a small path, the change in pressure is

$$dp = \frac{\partial p}{\partial r} dr + \frac{\partial p}{\partial h} dh = \rho\omega^2 r dr - \rho g dh.$$

Integrate this differential along the following integration path



to find

$$\begin{aligned} \int dp &= \int_0^r \rho\omega^2 r dr - \int_{h_c}^h \rho g dh \\ p - p_a &= \frac{1}{2}\rho\omega^2 r^2 - \rho g(h - h_c). \end{aligned}$$

c. At the fluid surface,  $p = p_a$ , so

$$0 = \frac{1}{2}\rho\omega^2 r^2 - \rho g(h - h_c) = \frac{1}{2}\rho\omega^2 r^2 - \rho g(y(r))$$

whence

$$y(r) = \frac{\omega^2}{2g} r^2.$$

Note that the profile  $y(r)$  is independent of  $\rho$ : water and mercury will have identical surface profiles.